



Department of Health

ANDREW M. CUOMO
Governor

HOWARD A. ZUCKER, M.D., J.D.
Commissioner

LISA J. PINO, M.A., J.D.
Executive Deputy Commissioner

Guidance for The New York State COVID-19 Vaccination Program May 21, 2021

Purpose and Background:

All individuals **12 years of age and older** that reside in the United States are eligible to be vaccinated. See Appendix A for guidance regarding necessary consent for individuals under 18 years of age.

Performance, throughput, effort, and effective administration of vaccines by providers continue to be key factors in making future vaccine allocations, along with equity and access.

All vaccine providers in New York State, including those located in the City of New York and those participating in federal programs, must follow New York State Department of Health (NYSDOH) guidance and directives, including the requirement to accurately and completely report doses administered to the appropriate immunization information system (NYSIIS or CIR) within 24 hours of vaccine administration per Executive Order 202.82 as extended by 202.89, and must maintain up-to-date inventory in such system.

Accurate and timely reporting to NYSIIS/CIR is critical, as this information can be used to allow individuals to display proof of vaccination, such as the Excelsior Pass.

Effective May 13, 2021, reporting to the NYS COVID-Vaccine Tracker is no longer required.

Eligible individuals:

All individuals 12 years of age and older that reside in the United States are eligible to be vaccinated.

On May 12, following the New York State Clinical Advisory Task Force's recommendation and CDC's adoption of the Advisory Committee on Immunization Practices' (ACIP) endorsement in response to the U.S. Food and Drug Administration's expansion of the [emergency use authorization \(EUA\) for the Pfizer-BioNTech COVID-19 Vaccine](#) to include adolescents 12 through 15 years of age, Governor Cuomo authorized all providers enrolled in the NYS COVID-19 vaccination program to expand eligibility for the Pfizer COVID-19 vaccine to 12 through 15-year olds, effective immediately.

Minors must present identification to verify that they are at least 12 years of age or have a parent/guardian attest on their behalf. See Appendix A for guidance regarding necessary consent for individuals under 18 years of age.

The mandatory [New York State COVID-19 Vaccine Form](#) includes demographic questions and a self-attestation and must be completed by all individuals prior to vaccination.

Updates to Pfizer-BioNTech COVID-19 Vaccine:

On May 19, the Food and Drug Administration (FDA) amended the Pfizer EUA to allow undiluted, thawed Pfizer-BioNTech COVID-19 Vaccine vials to be **stored in the refrigerator at 2°C to 8°C (35°F to 46°F) for up to one month**. Previously, thawed, undiluted vaccine vials could be stored in the refrigerator for up to five days. The updated storage and handling summary allowances for Pfizer are as follows:

- Ultra-cold (-80° to -60° C): May be stored in ultra-cold freezer until expiration or temporarily in the shipper with dry ice replenishment every five days for up to 30 days.
- Frozen (-25° to -15° C): May be stored in standard freezer up to 2 weeks.
- Refrigerated (2° to 8° C): Undiluted for up to 1 month. Cannot be refrozen once thawed/refrigerated.
- **Total storage time for Pfizer in freezer and refrigerator should not exceed 45 days (14 days frozen plus 31 days refrigerated). Beyond use dates for standard freezer/refrigeration must not exceed the labeled expiration date (i.e., not cumulative)**
- Punctured vials: At refrigerated or room temperature up to 6 hours

In addition, Pfizer COVID-19 Vaccine will now have a new, smaller packaging quantity of 450 doses available for allocation [beginning May 27](#), with earliest delivery date of [June 1](#). The 450-dose ordering package (NDC 59267-1000-03) will consist of three trays of 25 vials (150 doses) each and will ship ultra-frozen. Dry ice will NOT be sent in follow-up to the Pfizer 450-dose orders. Ancillary supply kit contents will be the same, with quantities adjusted for the smaller package. This new package size, combined with the updated storage requirements to allow Pfizer vaccine to be stored refrigerated for one month, means that vaccination providers without ultra-cold storage may now be able to split one order to support both first dose and second dose administration. The 1170-dose quantity of Pfizer vaccine will continue to be available as well and will still come with dry ice recharge 24 hours later if you have not opted out of dry ice shipments. Because only 35% of the U.S. Pfizer supply as of [June 1](#) will be in the 450-dose package size, we encourage providers with ultra-cold storage to continue to order the full 1170-dose package to enable those that do not have ultra-cold freezers for long-term storage to receive the smaller package size.

No Minimum Interval Between COVID-19 Vaccine and Other Vaccines:

On May 14, the CDC updated its [“Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States,”](#) to recommend that “COVID-19 vaccines and other vaccines **may now be administered without regard to timing**. This includes simultaneous administration of COVID-19 vaccines and other vaccines on the same day, as well as coadministration within 14 days.” Although COVID-19 vaccines were previously recommended to be administered a minimum of 14 days before or after other vaccines, that previous recommendation was out of an abundance of caution and not due to any known safety or immunogenicity concerns, and is no longer in effect. When deciding whether to co-administer another vaccine(s) with COVID-19 vaccines, providers should consider whether the patient is behind or at risk of becoming behind on recommended vaccines, their risk of vaccine-preventable disease (e.g., during an outbreak or occupational exposures), and the reactogenicity profile of the vaccines.

Vaccine Provider Responsibilities:

- Vaccine can be redistributed to another facility, provider, practice, or local health department that is enrolled in the COVID-19 vaccination program, with prior notice to the NYSDOH. To redistribute vaccine, facilities must submit a [completed redistribution form](#) to COVIDVaccineRedistribution@health.ny.gov and can proceed with the redistribution once submitted.
 - A provider may transport vaccine to another location for the purpose of holding a limited duration vaccination clinic without prior notification to the NYSDOH. If the provider is administering the doses and reporting doses administered against their own inventory in NYSIIS, all unused vaccine must be transported back to the original location at the conclusion of the clinic that day. The provider must retain possession and control of the vaccine for the duration of the transport and administration.
- All vaccine providers should minimize the amount of vaccine that goes unused, consistent with CDC guidance, which states that while enrolled providers must continue to follow best practices to use every dose possible, it should not be at the expense of missing an opportunity to vaccinate every eligible person when they are ready to get vaccinated. (See page 4 for further guidance.)
- Providers should not prefill more syringes than they can use within one hour. Prefilled syringes of Pfizer-BioNTech and Moderna vaccines must be used within six hours of filling; Janssen (Johnson & Johnson) vaccine must be used within two hours of filling. Excess prefilling can lead to waste if a clinic must end early or an excessive number of recipients fail medical screening or do not show up for their appointment. Please see [Guidance on Use of COVID-19 Vaccine Doses Remaining at End of Day or Clinic for Providers Participating in the New York State COVID-19 Vaccination Program](#) for more information.
- All facilities or practices are required to track vaccine uptake among their staff and must furnish uptake data to the NYSDOH via HERDS survey, or as directed by your agency or organization.

Each provider that receives vaccine:

- MUST ensure that each individual they vaccinate displays evidence of completed [NYS COVID-19 Vaccine Form](#) and attestation.
- Must make best efforts to use all vaccine doses within seven days of receipt by rapidly administering it to eligible individuals.
- All vaccine administered must be accurately and completely reported, using the New York State Immunization Information System (NYSIIS) or the Citywide Immunization Registry (CIR) in New York City, within 24 hours of administration and providers must maintain up-to-date inventory in such system. This is critically important to allow an individual to prove vaccination status.
 - Reporting to the NYS COVID-Vaccine Tracker is **no longer required**.
- With respect to **pharmacies**, pharmacists are authorized to vaccinate individuals 12 years of age and older for COVID-19, pursuant to the Seventh Amendment to Declaration Under the Public Readiness and Emergency Preparedness [PREP] Act for Medical Countermeasures Against COVID-19.

Message for COVID-19 Vaccine Clinical Trial Sites:

As a reminder, all COVID-19 vaccines administered in the State of New York must be entered in to NYSIIS or CIR. This includes any doses administered as part of an experimental arm of a clinical trial as well as doses offered and administered to participants in the control group (originally received placebo) after the clinical trial ended or at other time points per trial protocol. Staff at the participating site of the clinical trial must provide participants with a vaccination card and enter participant's immunization history into NYSIIS/CIR as applicable. Please note

that only vaccines that have been issued an Emergency Use Authorization or that have been approved by the United States Food and Drug Administration (FDA) can be entered.

The Second COVID-19 Vaccine Dose:

Pfizer-BioNTech and Moderna vaccines require two doses, whereas Janssen (Johnson & Johnson) vaccine requires only a single dose. The second dose must be administered 21 days (Pfizer-BioNTech vaccine) or 28 days (Moderna vaccine) after the first dose. To facilitate this, all providers **must** schedule the second dose appointment for recipients **at the time the first dose is administered**.

Individuals must receive two doses of the same vaccine (e.g., you must receive two doses of the Pfizer-BioNTech vaccine or two doses of the Moderna vaccine). They are **not** interchangeable. Please see [Guidance for Administration of the Second Dose of COVID-19 Vaccine](#) for additional information regarding administration of the second dose.

If an individual requests a second dose after missing the 42-day window, they should still be administered a second dose. There is no need to restart the series, pursuant to CDC guidance. Providers who have insufficient vaccine to administer a second dose that was delayed beyond the 42-day window should work with their Lead Hub Hospital.

Circumstances may arise where individuals need to receive their second dose at a different location than their first. Providers who have determined that the individual cannot return to the location where they received their first dose should schedule a second dose for these individuals or coordinate with the Regional Hub Lead Hospital to find a provider who has extra second doses of the appropriate vaccine to vaccinate the individual. Vaccine availability can also be located using the [CDC Vaccine Finder](#). Individuals should not be tasked with locating second dose appointments. This obligation is on the provider who administered the first dose.

Special Considerations for Individuals Receiving Their First Dose Outside New York State:

Individuals who received their first dose of COVID-19 vaccine outside of New York State will not have a record of this dose in NYSIIS or CIR. Providers administering a second dose should either enter the first dose in NYSIIS/CIR as part of the historical record using data listed on the individual's COVID-19 Vaccination Record Card OR advise the patient that they should ask their primary care provider to enter their first dose into NYSIIS/CIR so the state has a full record of both doses of COVID-19 vaccine.

Universal Doses:

Effective May 11th, New York State moved to a "Universal Dose" administration process for all multi-dose COVID-19 vaccine types. All doses are now considered universal doses, which means that doses can be used as either a first dose or a second dose, regardless if they were originally shipped to providers as a first dose or a second dose. This does NOT eliminate the obligation of the provider to schedule second dose appointments at the time the first dose is administered.

First and second doses may also be drawn interchangeably from the same vial. With all doses considered a universal dose, please utilize a first in, first out rule to manage inventory. This includes storing newly received vaccine in the freezer until it is needed. **COVID-19 vaccine providers should continue to follow their jurisdiction's (NYC or NYS) vaccine ordering and inventory guidance to request their weekly vaccine allocations.**

Extra Doses of Pfizer-BioNTech and Moderna:

Vials of both Pfizer-BioNTech and Moderna may contain at least one extra dose of vaccine. Depending on the type of needle and syringes used, additional vaccine may remain in the vial. Vaccine administrators may use any extra vaccine that can be easily drawn up in a syringe to meet the full dose requirements. Extra vaccine fluid from more than one vial **CANNOT** be combined to produce extra doses.

This is particularly important because the vaccination does not contain preservatives. Enter all vaccines given into NYSIIS/CIR, including any additional vaccines given, however do not modify inventory in anticipation of extra doses. For additional information please see [Pfizer-BioNTech](#) guidance and [Moderna](#) guidance for extra doses. Extra vaccine has not been observed in the Janssen (Johnson & Johnson) vials beyond the expected five doses.

Responsible Wastage

The CDC released guidance on May 11th regarding wastage with the critical message to “take every opportunity to vaccinate every eligible person.” As more vaccination opportunities are created, the likelihood of leaving unused doses in a vial may increase. While enrolled providers must continue to follow best practices to use every dose possible, it should not be at the expense of missing an opportunity to vaccinate every eligible person when they are ready to get vaccinated.

To ensure providers do not miss an opportunity to vaccinate every eligible person:

- Providers must follow [clinical best practice for vaccination as well as best practices when managing inventory](#) to maximize vaccination and minimize dose wastage.
- Providers should not miss any opportunities to vaccinate every eligible person who presents at a vaccination site
 - Consider establishing and promoting standing vaccination days or half-days to increase likelihood of larger numbers of people presenting for vaccination on the same day.
 - Vaccinate family members or friends who accompany patients to medical visits even if they are not established patients at the vaccinating practice.
 - Continue outreach to employers or other community partners that have a large membership or network to arrange vaccination events.
 - As contingency plan, vaccine providers should attempt to contact additional persons (i.e., from a standby list or through personal contacts of persons being vaccinated) to use as many vaccine doses as possible.
 - Once punctured, multidose vials must be used within:
 - 12 hours (Moderna)
 - 6 hours (Pfizer)
 - 6 hours (refrigerated) or up to 2 hours at room temperature (J&J/Janssen)

Mandatory Vaccine Form:

All individuals receiving the COVID-19 vaccine **must** complete the [New York State COVID-19 Vaccine Form](#) for the first dose, and attest that they are eligible to be vaccinated. All practices, providers, and entities must confirm adherence to this requirement at the time of vaccine administration.

Vaccine Safety:

Post-vaccination monitoring is an essential part of the COVID-19 vaccination program. The Centers for Disease Control and Prevention (CDC) is promoting and encouraging all those being vaccinated to participate in V-Safe, a smart-phone based application that will allow those vaccinated to enter their symptoms in the days after vaccination using text messaging. V-Safe also provides reminders for the second dose and telephone follow up for anyone who reports medically significant adverse events. V-Safe materials can be found at <http://www.cdc.gov/vsafe>, including a V-Safe information sheet. Please print out the information sheet and hand to each person vaccinated. You must report any adverse events that occur after vaccination to the Vaccine Adverse Events Reporting System (VAERS) at info@VAERS.org or by calling 1-800-822-7967.

Equity and Access:

Effort must be made to do outreach to persons 12 years of age and older in all communities and settings. Persons in areas that have a high social vulnerability index are particularly vulnerable to COVID-19 and should be notified about how they can receive vaccine. Every effort should be made to increase their access to vaccination opportunities.

Communicating the Plan:

Please be sure to clearly communicate this critical guidance to all staff involved in the vaccination program.

This guidance is in effect from the date of issuance until it is updated, or additional guidance is issued by NYSDOH. For questions, please contact the New York State Department of Health, Bureau of Immunization at COVID19vaccine@health.ny.gov.

New York State Vaccination Program Guidance Appendix A – Minor Consent

All individuals 12 years of age and older that reside in the United States are eligible to be vaccinated. **However, minors 12 through 17 are NOT authorized to receive the Janssen/Johnson & Johnson or Moderna COVID-19 vaccines. They may ONLY receive Pfizer at this time pursuant to the FDA Emergency Use Authorization (EUA). Children under 12 years of age are not yet authorized to receive ANY COVID-19 vaccine.**

It is important to verify the age of individuals who appear to be a minor to confirm eligibility and ensure the administration of the proper COVID-19 vaccine.

Proof of age should be requested but is not required where the parent or guardian is available to attest to the minor's age. Documentary proof may include (but is not limited to):

- Driver's license or non-driver ID;
- Birth certificate issued by a state or local government
- Consulate ID
- Current U.S passport or valid foreign passport
- Permanent resident card
- Certificate of Naturalization or Citizenship
- Life insurance policy with birthdate
- Parent/Guardian attestation

Minor Consent

16 and 17-year olds

For all minors, a parent or legal guardian must provide consent for vaccination. For minors 16 or 17 years of age, such consent should be provided either in person or by phone, at the time of vaccine appointment. Providers may elect whether to accept a written statement of consent from the parent or guardian, where the parent or guardian is not available by phone to provide consent to vaccinate an unaccompanied minor. The [NYS COVID-19 Immunization Screening and Consent Form](#) may be considered for this purpose.

12 through 15-year olds

For minors who are 12 through 15 years of age, additionally, an adult caregiver should accompany the minor. If the adult caregiver is not the parent/guardian, the adult caregiver should be designated by the parent/guardian. The parent/guardian must still provide consent to the vaccination.