



Information for Health Care Professionals about the Screening Checklist for Administration of the Pfizer-BioNTech COVID-19 Vaccine for Children Ages 5-11

Updated: April 1, 2022

Note: For summary information on contraindications and precautions to vaccines, go to the ACIP's General Best Practice Guidelines for Immunization at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html.

1. Are you between the ages of 5 and 11 years old?

If yes, proceed with screening question #3. If no, advise the parent or guardian that this patient is not currently eligible for the 5-11 years old pediatric Pfizer BioNTech Covid-19 formulation (10 µg/ 0.2 mL). Instead, this patient may receive the Pfizer BioNTech Covid-19 adult formulation for children 12 years old or older (30 µg/ 0.3 mL). There is no COVID-19 vaccine currently authorized for a child less than 5 years old.

If a child turns from 11 to 12 years of age in between their first and second dose in the primary regimen, they may receive, for either dose, either: (1) the Pfizer-BioNTech COVID-19 Vaccine formulation for children aged 5–11 years (each 0.2 ml dose containing 10 µg in an orange cap vial); or (2) COMIRNATY or the Pfizer-BioNTech COVID-19 Vaccine formulation authorized for use in individuals 12 years of age and older (each 0.3 mL dose containing 30 µg in a purple cap vial). If they receive the 10 µg dose for their second dose instead of the 30 µg dose, this is not considered an error, VAERS reporting is not required, and the child is considered fully vaccinated. However, based on clinical judgment, a repeat dose of the 30µg adult formulation may be administered 21 days after the second pediatric formulation dose was administered.

2. Are you 12 years old or older?

If the patient is 12 years old, or greater, they are eligible for the adult Pfizer vaccine formulation. "Information for Health Care Professionals about the Screening Checklist for the COVID-19 Vaccine" for individuals 12 years old or older can be found on the [New York State COVID-19 Vaccine Information for Providers](#) page.

3. Are you feeling sick today?

If yes, refer to the vaccination site healthcare provider for assessment of current health status. If patient is feeling moderately or severely ill, do not vaccinate at this time. Ask the parent or guardian to return with the patient when symptoms improve.

4. In the last 10 days, have you had a COVID-19 test because you had symptoms and are still awaiting your test results or been told by a health care provider or health department to isolate or quarantine at home due to COVID-19 infection or exposure?

- If yes, advise the parent or guardian that the patient must return to isolation or quarantine and reschedule for after isolation/quarantine ends.
- If the patient was diagnosed with COVID-19 greater than 10 days ago and has been asymptomatic for 72 hours or more, patient may be vaccinated.

- If the patient has had a test in the last 10 days, ask for the result. If positive, send them home. If negative, they can proceed to vaccination. If the result is unsure or unknown, advise the patient to return once a negative test has been confirmed or 10 days have passed since a positive test.
- Persons with a history of multisystem inflammatory syndrome in children (MIS-C) or in adults (MIS-A) should consider delaying vaccination until they have recovered from their illness and for 90 days after the diagnosis of MIS-C or MIS-A. However, the parent or guardian can choose to have the patient vaccinated. For further information on counseling a parent or guardian with a child who has a history of MIS-C or MIS-A regarding COVID-19 vaccines, please see the Centers for Disease Control and Prevention’s (CDC) section on MIS-C and MIS-A in their “Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States” available at: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html>.

5. Have you been treated with antibody therapy or convalescent plasma for COVID-19 in the past 90 days (3 months)? If yes, when did you receive the last dose?

If yes, notify them that they should defer future doses of Evusheld™ for at least 2 weeks after vaccination. However, COVID-19 vaccination need not be delayed due to previous antibody therapy nor receipt of convalescent plasma.

6. Have you ever had an immediate allergic reaction, such as hives, facial swelling, difficulty breathing, anaphylaxis to any vaccine, injection, or shot or to any component of the COVID-19 vaccine, or a severe allergic reaction (anaphylaxis) to anything?

If yes, then refer to the vaccination site healthcare provider for assessment of allergic reaction. Review the ingredient lists at: https://www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fvaccines%2Fcovid-19%2Finfo-by-product%2Fclinical-considerations.html#Appendix-C.

Contraindications to COVID-19 vaccine:

- Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine.
- Known diagnosed allergy to a component of the COVID-19 vaccine
- People with a contraindication to one of the mRNA COVID-19 vaccines should not receive a dose of the Pfizer COVID-19 vaccine.

Precautions to COVID-19 vaccine: (Refer to your organization’s protocol to see whether individuals with a precaution to vaccination warrant further evaluation.)

- Immediate (onset within 4 hours after vaccination), but non-severe, allergic reaction after a previous dose of COVID-19 vaccine
- Immediate allergic reaction to any other vaccine or injectable therapy (i.e., intramuscular, intravenous, or subcutaneous vaccines or therapies excluding subcutaneous immunotherapy for allergies).
- Individuals with a contraindication to one type of COVID-19 vaccine, or a component of a COVID-19 vaccine may have a precaution to the other.
 - Consultation with an allergist-immunologist should be considered to help determine if the patient can safely receive vaccination, and vaccination of these individuals should only be undertaken in an appropriate clinical setting under the supervision of a healthcare provider experienced in the management of severe allergic reactions.
 - Note: These individuals should not be administered COVID-19 vaccine at State-operated vaccination sites.

- For mRNA COVID-19 vaccines, for history of myocarditis or pericarditis after a dose of an mRNA COVID-19 vaccine, a subsequent dose of COVID-19 vaccine should generally be avoided (there is a separate question on the screening and consent form regarding myocarditis and pericarditis – question #9).

For patients who are determined eligible for COVID-19 vaccination after assessment of allergy history, a 30-minute post-vaccination observation period is needed for the following:

- Patients with a history of an immediate allergic reaction of any severity to any other vaccine or injectable therapy.
- Patients with a contraindication to any type of COVID-19 vaccine (e.g., mRNA or Janssen viralvector).
- Patients with a history of anaphylaxis due to any cause.

7. Do you have cancer, leukemia, HIV/AIDS, or any other condition that weakens the immune system?

If yes, ask the parent or guardian if they would like to have a discussion with the vaccination site healthcare provider about what is known and not yet known about COVID-19 vaccine for immunocompromised people. You can tell the parent or guardian that the patient may have a less strong immune response to the vaccine but may still get vaccinated. Patient may be vaccinated if the parent or guardian chooses to, and they are not required to go to medical evaluation.

8. Do you take any medications that affect your immune system, such as cortisone, prednisone or other steroids, anticancer drugs, or have you had any radiation treatments?

If yes, ask the parent or guardian if they would like to have a discussion with the vaccination site healthcare provider about what is known and not yet known about COVID-19 vaccine for immunosuppressed people. You can tell the parent or guardian that they may have a less strong immune response to the vaccine but may still get vaccinated. Patient may be vaccinated if the parent or guardian chooses to, and they are not required to go to medical evaluation.

9. Do you have a bleeding disorder, a history of blood clots or are you taking a blood thinner?

If yes, refer to parent or guardian to their healthcare provider to assess the patient’s bleeding risk and thrombosis history. Persons with a history of Thrombosis with Thrombocytopenia Syndrome (TTS) following receipt of Janssen COVID-19 vaccine or of any other immune-mediated thrombosis and thrombocytopenia, such as Heparin-Induced Thrombocytopenia (HIT) within the past 90 days should be offered an mRNA COVID-19 vaccine (i.e., Pfizer vaccine). If a person with a bleeding disorder or taking a blood thinner is cleared for vaccination, then administer vaccine using a 23-gauge or smaller caliber needle and apply firm pressure on the site of vaccination, without rubbing, for at least 2 minutes after vaccination.

10. Do you have a history of myocarditis (inflammation of the heart muscle) or pericarditis (inflammation of the lining around the heart)?

If yes:

- Evaluate if this history was in relation to a dose of mRNA vaccine. If it was not, then the patient can receive any U.S. Food and Drug Administration (FDA) authorized COVID-19 vaccine after complete resolution of a myocarditis or pericarditis episode.
- Until additional safety data are available, experts advise that people who develop myocarditis or pericarditis after a dose of an mRNA COVID-19 vaccine generally **should not** receive a subsequent dose of any COVID-19 vaccine. If after a risk assessment, the decision is made to receive a subsequent COVID-19 vaccine dose, the person should wait until at least after their episode of myocarditis or pericarditis has resolved (i.e., resolution of symptoms, no evidence of ongoing heart inflammation or sequelae as determined by patient’s clinical team). Decisions to proceed with vaccination should include conversations with the patient, parent/legal representative, and the

clinical team, including a cardiologist. Considerations for vaccination may include:

- The myocarditis or pericarditis was considered unrelated to mRNA COVID-19 vaccination (e.g., due to SARS-CoV-2 or other viruses), especially if the myocarditis or pericarditis diagnosis occurred more than 3 weeks after the most recent COVID-19 vaccine.
- Personal risk of severe acute COVID-19 disease (e.g., age, underlying conditions).
- Level of COVID-19 community transmission and personal risk of infection.
- Timing of immunomodulatory therapeutics; ACIP's [general best practice guidelines for immunization](#) can be consulted for more information.
- For the full CDC interim clinical considerations regarding a history of myocarditis and/or pericarditis, please see the CDC's [COVID-19 Vaccines Currently Authorized in the United States](#) and [Clinical Considerations: Myocarditis and Pericarditis after Receipt of mRNA COVID-19 Vaccines Among Adolescents and Young Adults](#).

11. Do you have a history of MIS-C (Multisystem Inflammatory Syndrome in Children)?

If yes, and they have not yet received vaccine, they should defer COVID-19 vaccines until clinical recovery from MIS-C, including return to normal cardiac function, plus at least 90 days since their MIS-C diagnosis.

For people who had MIS-C fewer than 90 days after the date of their most recent COVID-19 vaccine dose, further COVID-19 vaccine doses should be deferred until additional data are available.

12. Have you received a previous dose of the Pfizer, Moderna or Janssen COVID-19 vaccine?

If yes, verify if this the patient is still under the age of 12 years old. Children ages 5-11 who already received a first dose of a two-dose mRNA COVID-19 vaccine should be offered the age-appropriate second dose of an mRNA vaccine, which at this time is only the Pfizer-BioNTech COVID-19 Vaccine formulation for children ages 5-11 (10 µg).

If the patient is 12 years old or older, please counsel the parent or guardian on current guidelines to provide a Pfizer COVID-19 vaccine adult formulation (30 µg) to anyone in the age range. The Moderna and Janssen vaccines are only authorized for those 18 years or older.

- Have you received a previous dose of a COVID-19 vaccine recognized by the WHO but NOT by the FDA (AstraZeneca - VAXZEVRIA, Sinovac - CORONAVAC, Serum Institute of India - COVISHIELD, Sinopharm / BIBP, Covaxin, Serum Institute of India - Covovax / Novavax- NUVAXOVID)? If yes, identify if the patient has received a complete or partial series of the vaccine. If the patient received a complete series (e.g., 2 doses), CDC considers them to be fully vaccinated and no additional doses are needed.
- If a patient received a partial series of a WHO authorized COVID-19 vaccine that is not currently authorized for use in the U.S. by the FDA, the CDC does NOT consider these persons to be fully vaccinated. They should be offered a two-dose series of the age-appropriate COVID-19 vaccine (i.e., Pfizer-BioNTech COVID-19 Vaccine formulation for children ages 5-11).
- If the patient received either a partial series or complete series of a COVID-19 vaccine that is not authorized for use by either the WHO or the FDA, the CDC does NOT consider these persons to be fully vaccinated. They should be offered a two-dose series of the age-appropriate COVID-19 vaccine (i.e., Pfizer-BioNTech COVID-19 vaccine formulation for children ages 5-11).

*** Anyone answering "Unknown" to any screening question should be referred to the medical director or responsible healthcare provider at the POD or clinic to further assess their answer to that question (e.g., the person might not have understood the question and the healthcare provider could explain it further).**